



Gata5 is required for the development of the heart and endoderm in zebrafish

Jeremy F. Reiter, Jonathan Alexander, Adam Rodaway, Deborah Yelon, Roger Patient, Nigel Holder and Didier Y.R. Stainier

Genes & Dev. 1999 13: 2983-2995

Access the most recent version at doi:[10.1101/gad.13.22.2983](https://doi.org/10.1101/gad.13.22.2983)

References

This article cites 57 articles, 33 of which can be accessed free at:
<http://www.genesdev.org/cgi/content/full/13/22/2983#References>

Article cited in:
<http://www.genesdev.org/cgi/content/full/13/22/2983#otherarticles>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#)

Notes

To subscribe to *Genes and Development* go to:
<http://www.genesdev.org/subscriptions/>

Gata5 is required for the development of the heart and endoderm in zebrafish

Jeremy F. Reiter,¹ Jonathan Alexander,¹ Adam Rodaway,² Deborah Yelon,¹ Roger Patient,² Nigel Holder,^{3,4} and Didier Y.R. Stainier^{1,5}

¹Department of Biochemistry and Biophysics, Programs in Human Genetics and Developmental Biology, University of California at San Francisco, San Francisco, California 94143-0448 USA; ²Developmental Biology Research Centre, The Randall Institute, King's College London, London WC2B 5RL, UK; ³Department of Anatomy and Developmental Biology, University College London, London WC1E 6BT, UK

The mechanisms regulating vertebrate heart and endoderm development have recently become the focus of intense study. Here we present evidence from both loss- and gain-of-function experiments that the zinc finger transcription factor Gata5 is an essential regulator of multiple aspects of heart and endoderm development. We demonstrate that zebrafish Gata5 is encoded by the *faust* locus. Analysis of *faust* mutants indicates that early in embryogenesis Gata5 is required for the production of normal numbers of developing myocardial precursors and the expression of normal levels of several myocardial genes including *nkx2.5*. Later, Gata5 is necessary for the elaboration of ventricular tissue. We further demonstrate that Gata5 is required for the migration of the cardiac primordia to the embryonic midline and for endodermal morphogenesis. Significantly, overexpression of *gata5* induces the ectopic expression of several myocardial genes including *nkx2.5* and can produce ectopic foci of beating myocardial tissue. Together, these results implicate zebrafish Gata5 in controlling the growth, morphogenesis, and differentiation of the heart and endoderm and indicate that Gata5 regulates the expression of the early myocardial gene *nkx2.5*.

[Key Words: Gata; cardia bifida; *nkx2.5*; organogenesis; morphogenesis; liver]

Received July 22, 1999; revised version accepted October 5, 1999.

Fusion of the cardiac primordia is the first in a complex series of morphogenetic steps required to form a functional heart. The cardiac primordia are contained within the anterior lateral plate mesoderm (ALPM). During segmentation stages, the bilateral populations of ALPM migrate medially and fuse to form a cell layer that crosses the embryonic midline. As part of this process, the two cardiac primordia meet and fuse at the embryonic midline to form the definitive heart tube. In the absence of migration, the two cardiac primordia develop in their original lateral positions, resulting in a phenotype called cardia bifida. Six zebrafish mutations causing cardia bifida were identified in the large-scale screens for genes regulating embryonic development: *faust* (*fau*), *casanova* (*cas*), *one-eyed pinhead* (*oep*), *bonnie and clyde*, *miles apart*, and *natter* (Chen et al. 1996; Stainier et al. 1996).

Two of these mutations, *oep* and *cas*, also disrupt endoderm formation (Schier et al. 1997; Alexander et al. 1999), underscoring the developmental relationship between the heart and endoderm. Classical embryological experiments initially identified the ability of endoderm to induce rhythmic contraction in cardiac explants (Jacobson and Sater 1988). More recent experiments have

confirmed that the endoderm is required for avian cardiac myofibrillogenesis (Gannon and Bader 1995). The endoderm is also implicated in the earlier process of pre-cardiac mesoderm induction; coculture of *Xenopus* deep endoderm with the heart primordia enhances cardiogenesis (Nascone and Mercola 1995), and BMP2, which is secreted by the anterior endoderm, is sufficient to induce expression of myocardial genes in avian noncardiogenic mesoderm (Schultheiss et al. 1997).

Heart and endoderm development in the mouse both require the zinc finger transcription factor Gata4 (Kuo et al. 1997; Molkentin et al. 1997). Although 35% of *gata4* null embryos arrest at the egg cylinder stage for unknown reasons, those that do gastrulate fail to form a fused heart or a foregut and develop partially outside the yolk sac and amnion. Gene expression within the pre-cardiac mesoderm is not significantly affected, suggesting that cardiac differentiation is not the primary defect. In support of this conclusion, chimera analyses indicate that Gata4 function in the visceral and/or anterior definitive endoderm restores ventral closure and cardiac fusion to otherwise *gata4* null embryos (Narita et al. 1997).

Gata4 is one of a family of six vertebrate Gata transcription factors, each of which contains two zinc fingers and binds to regulatory elements containing A/TGATAA/G or closely related sequences. The *gata*

⁴Deceased.

⁵Corresponding author.

E-MAIL didier_stainier@biochem.ucsf.edu; FAX (415) 476-3892.

genes can be divided into two groups on the basis of sequence homology and expression pattern. The members of the first group, *gata1*, *gata2*, and *gata3* have unique roles in hematopoiesis (for review, see Orkin and Zon 1997), whereas *gata4*, *gata5*, and *gata6* are expressed in the ALPM and endoderm (for review, see Evans 1997). Studies of *cis*-regulatory elements have suggested that *gata4*, *gata5*, and *gata6* have important roles in promoting the expression of both myocardial and endodermal genes. Transfection of heterologous cells with Gata factors activates transcription of myocardial genes including α -myosin heavy chain (α -MHC), cardiac troponin C, and B-type natriuretic peptide (BNP) (for review, see Evans 1997), as well as endodermal genes such as *IFABP* and *HNF4* (Gao et al. 1998; Morrissey et al. 1998). Evidence that Gata factors directly activate the transcription of myocardial and endodermal genes has come from studies identifying functionally important Gata-binding sites in the regulatory regions of the myocardial genes α -MHC and BNP (for review, see Evans 1997) and the liver enhancer of *serum albumin* (Bossard and Zaret 1998). Recently, Gata-binding sites have been identified in two different regulatory elements of *nkx2.5*, one of the earliest known markers of the myocardial precursors (Searcy et al. 1998; Lien et al. 1999). Each of these elements activates expression in a manner similar to the endogenous *nkx2.5* promoter and depends on Gata sites for activity.

Experiments in chick and *Xenopus* provide further in vivo evidence for the role of *gata* factors in heart development. Inhibition of Gata activity through the addition of three antisense oligonucleotides complementary to *gata4*, *gata5*, and *gata6* produces variable abnormalities in avian cardiac morphogenesis (Jiang et al. 1998). Ectopic expression of *gata4* and *gata5* in *Xenopus* embryos leads to the premature expression of the myocardial genes α -cardiac actin and α -MHC (Jiang and Evans 1996). In contrast, sustaining the expression of *Xenopus gata6* after it has normally declined inhibits expression of markers of terminal myocardial differentiation, perhaps by maintaining the precardiac mesoderm in an undifferentiated state (Gove et al. 1997).

gata genes also play evolutionarily conserved roles in endoderm development, as the *Drosophila gata* gene *serpent* and the *Caenorhabditis elegans gata* gene *elt-2* are essential for differentiation of the embryonic gut (Rehorn et al. 1996; Fukushige et al. 1998). Another *C. elegans gata* gene, *end-1*, is implicated in the earlier process of endoderm specification (Zhu et al. 1997, 1998).

Here, we show that the zebrafish *fau* locus is essential for multiple aspects of heart and endoderm development and that *fau* encodes Gata5. Analysis of *fau* mutants reveals that Gata5 is required for the normal expression of *nkx2.5* and for other aspects of myocardial differentiation. Mutations in *gata5* also prevent the formation of normal amounts of cardiac and endodermal tissue and inhibit the normal morphogenesis of the heart, pharyngeal endoderm, and gut. Furthermore, we demonstrate that the overexpression of *gata5* is sufficient to activate ectopic expression of *nkx2.5* and other cardiac genes and

to cause the formation of ectopic regions of rhythmically contracting tissue. Thus, our results establish Gata5 as a critical regulator of *nkx2.5* expression and of multiple aspects of cardiac and endoderm development.

Results

fau mutants display defects in heart development

The zebrafish *fau* locus is defined by two alleles; *fau*^{tm236a} is a previously reported, ethylnitrosourea (ENU)-induced allele (Chen et al. 1996), whereas the second allele, *fau*^{s26}, arose spontaneously in an unrelated line. Both alleles segregate as fully penetrant, completely recessive mutations and cause equivalent phenotypes. The transheterozygous phenotype is also fully penetrant and is indistinguishable from the homozygous phenotypes.

Most conspicuously, *fau* mutations disrupt formation of the definitive heart tube, an early step in cardiac morphogenesis. Heart-tube formation depends upon the medial migration and fusion of the bilateral cardiac primordia (Stainier et al. 1993). In *fau* mutants this step of cardiac morphogenesis usually fails to occur, resulting in cardia bifida. However, 0%–45% of the *fau* mutant embryos in a given clutch produce fused, although morphologically abnormal, hearts. This variable expressivity is present in both alleles and also pertains to other aspects of the phenotype.

To examine when and how *fau* functions in heart development, we examined the expression of markers of the developing cardiac primordia. In *fau* mutants at 18.5 hr postfertilization (hpf), myocardial precursors expressing cardiac myosin light chain 2 (*cmlc2*) (Yelon et al. 1999) and endocardial precursors expressing *tie2* (Lyons et al. 1998) fail to migrate to the embryonic midline (Fig. 1B,C), demonstrating that the morphogenesis of both tissues that comprise the zebrafish heart is affected. Development of more posterior lateral plate mesodermal derivatives, such as the fin bud and blood, are unaffected in *fau* mutants (data not shown). Therefore, *fau* is essential for the proper morphogenesis of both of the cellular components of the zebrafish heart tube but is not generally required for the development of lateral plate mesodermal derivatives.

Mutations in *fau* lead to the abnormal expression of genes encoding myocardial sarcomere components. *cardiac troponin T*, *tropomyosin*, *cmlc1*, and *cmlc2* are all expressed in fewer cells in *fau* mutants than in wild-type siblings (Fig. 1B,D; data not shown). Detailed examination of myocardial sarcomere gene expression in *fau*^{tm236a} mutants reveals a differential requirement for the *fau* gene product. For example, *cmlc1* expression is more profoundly reduced in *fau*^{tm236a} mutants than is the expression of *cmlc2* (Fig. 1B,D, middle panels). This distinction is less apparent in *fau*^{s26} mutants as too few cells express sarcomere genes in these mutants to allow easy comparison. The more severe defect in *cmlc2* and *cmlc1* expression observed in *fau*^{s26} mutants than in *fau*^{tm236a} mutants indicates that the *fau*^{s26} allele is the stronger of the two.




Figure 1. *fau* mutants display abnormal cardiac morphogenesis and differentiation. Expression of *cmlc2* (B), *tie2* (C), *cmlc1* (D), *gata6* (E), *gata4* (F) and *nkx2.5* (G) in wild-type embryos and *fau* mutant siblings at the 19-somite (18.5 hpf; A,B,C,D), 12-somite (15 hpf; E,F) and 6-somite (12 hpf; G) stages. All views are dorsal with anterior to the top. (A) At the 19-somite stage, both *fau*^{tm236a} and *fau*^{s26} mutants are morphologically indistinguishable from wild-type siblings. The blue box approximately outlines the cardiogenic region, which lies ventral and just posterior to the midbrain–hindbrain boundary. B and D depict enlarged views of this region. (B) In wild-type embryos, the myocardial primordia express *cmlc2* and have fused at the embryonic midline (left). In contrast, the myocardial primordia of *fau*^{tm236a} mutants do not migrate to the embryonic midline, resulting in cardia bifida (middle). The myocardial primordia of *fau*^{s26} mutants also fail to migrate to the embryonic midline (right) and in addition display a severe reduction in the number of cells expressing *cmlc2*. (C) Formation of the endocardium (arrow) also depends on the migration of precursors from the ALPM to the embryonic midline. In *fau* mutants, endocardial precursors expressing *tie2* are absent from the midline. *tie2* expression in endothelial cells (arrowhead) reveals that more anterior vasculature forms normally in *fau* mutants. (D) *cmlc1* is also expressed in the myocardial primordia. In the right *fau*^{tm236a} cardiac primordium (middle), <20 cells express *cmlc1* whereas >120 cells express *cmlc1* in each wild-type primordium (left). In contrast, *fau*^{tm236a} mutants and wild-type siblings contain approximately equal numbers of *cmlc2*-expressing cells (see B). (E) *gata6* is widely expressed throughout the ALPM throughout somitogenesis. *gata6* expression is not perturbed in *fau* mutants. (F) In contrast, ALPM expression of *gata4* is slightly reduced in *fau* mutants. (G) *nkx2.5* expression is also reduced in *fau* mutants. Although the number of cells expressing *nkx2.5* is only moderately reduced in *fau*^{tm236a} mutants (middle), *nkx2.5* expression is severely diminished in *fau*^{s26} mutants (right).

During earlier segmentation stages before the migration of the cardiac primordia to the midline, the ALPM exists as two parallel stripes of cells lateral to the head and anterior notochord. *gata6*, a gene expressed throughout the ALPM, is expressed normally in *fau* mutants (Fig. 1E). However, the expression of the related gene *gata4* in the ALPM is slightly reduced (Fig. 1F). Together, these results indicate that *fau* is not required for the formation of the ALPM but is critical for the expression of cardiac genes within this tissue.

nkx2.5, an Nk homeobox gene required for heart development (for review, see Evans 1999), is also expressed in part of the ALPM from early in development. Expression of *nkx2.5* is dramatically reduced in *fau* mutants (Fig. 1G), demonstrating that *fau* is a direct or indirect regulator of *nkx2.5* expression and is required at a very early stage of ALPM differentiation. *fau*^{tm236a} mutants display less severely diminished *nkx2.5* expression (Fig. 1G), further indicating that *fau*^{s26} is the stronger allele.

Later in development, the zebrafish heart differentiates into an anterior ventricle and a posterior atrium. Immunofluorescent staining using monoclonal antibodies that recognize myosin heavy chain epitopes differentially expressed in ventricular and atrial tissues (Stainier and Fishman 1992) reveals that ventricular and atrial differentiation does not depend on cardiac fusion. For example, a robust amount of both ventricular and atrial

tissue forms in the cardia bifida mutant *cas* (Fig. 2A,B). However, *fau*^{tm236a} myocardia, in addition to being smaller than *cas* myocardia, contain proportionally much less ventricular tissue (Fig. 2B,C). This defect is also observable at earlier stages, before the cardiac primordia normally fuse. At the 18-somite stage the expression of *ventricular myosin heavy chain* (*vmhc*), a marker of ventricular myocardium (Yelon et al. 1999), is dramatically reduced in *fau*^{tm236a} mutants (Fig. 2D). *vmhc* expression is more severely reduced, or even absent, in *fau*^{s26} mutants (Fig. 2D).

fau mutants display defects in endoderm morphogenesis

As embryological experiments in chick and amphibia have suggested that the endoderm is a source of instructive signals for the precardiac mesoderm, we investigated the effects of the *fau* mutations on endoderm development. From 24 to 48 hpf, the zebrafish gut endoderm coalesces at the midline where it will form a tube, loop to the left, and differentiate into the liver, pancreas, swim bladder, and epithelium of the gastrointestinal tract (Warga and Nüsslein-Volhard 1999). The anterior gut endoderm expresses *axial*, a homolog of *HNF3β* (Strahle et al. 1996), whereas the posterior endoderm expresses *fkf2*, a homolog of *HNF3γ* (Odenthal and




Figure 2. Formation of ventricular myocardial tissue is abnormal in *fau* mutants. Mutants (33 hpf) and wild-type siblings were stained with MF20 and S46 antibodies (A–C). Red fluorescence indicates MF20 (TRITC) staining of ventricular myocardium and yellow fluorescence indicates the overlap of MF20 (TRITC) and S46 (FITC) staining of atrial myocardium. A–C show ventral views with anterior to the top. D shows dorsal views of *vmhc* expression at the 18-somite stage (18 hpf). (A) Wild-type embryos develop a single, looped heart patterned into a ventricle (red) and atrium (yellow). (B) *cas* mutants display cardia bifida and form both ventricular and atrial tissue. (C) *fau* mutants also display cardia bifida but form less total myocardium and only a very small amount of ventricular tissue. Here, one myocardium (left) has not formed any ventricular tissue. (D) *vmhc* is expressed in the preventricular component of the developing myocardium. *vmhc* expression is dramatically reduced in *fau* myocardial primordia even prior to cardiac fusion. *fau*^{s26} mutants display a variable loss of *vmhc* expression; (right) a *fau*^{s26} mutant in which less than five cells weakly express *vmhc* in the left ALPM (arrow), whereas none do on the right.

Nüsslein-Volhard 1998). In *fau* mutants, *axial*-expressing cells fail to coalesce at the midline, disrupting anterior gut tube morphogenesis (Fig. 3A, arrow). The posterior endoderm displays more variable defects that range from the near absence of gut to the failure of gut looping (Fig. 3B).

fau mutations also disrupt the morphogenesis of the pharyngeal endoderm. In *fau* mutants, the pharyngeal pouches are separated by a larger mediolateral distance (Fig. 3C), a defect that is reminiscent of the inability of the gut endoderm and precardiac mesoderm to migrate to the embryonic midline. In addition, the anterior pharyngeal pouches are dysmorphic in *fau*^{tm236a} mutants (Fig. 3C, arrow), whereas the most anterior *nkx2.3*-expressing cells appear to be absent in *fau*^{s26} mutants (Fig. 3D, arrow).

fau encodes *Gata5*

Initially, we mapped the *fau* locus 1.5 cM proximal to the simple sequence repeat marker z20895 on linkage group 23 (Shimoda et al. 1998). Using an enriched selection for amplified fragment length polymorphisms (AFLPs), we identified a number of markers located within a centimorgan of *fau*. In a parallel approach, we mapped *gata5*, a gene dynamically expressed in both the

endoderm and cardiac mesoderm (see below), to the same genomic region using the Goodfellow radiation hybrid panel (Kwok et al. 1998). Linkage analysis using a restric-





Figure 3. Endoderm development is abnormal in *fau* mutants. Expression of *axial* (A), *fkd2* (B), and *nkx2.3* (C,D) in wild-type embryos and *fau* mutant siblings at 44 hpf (A,B,D) and 28 hpf (C). All views are dorsal with anterior to the top. (A) *axial* is expressed in the anterior gut endoderm and the ventral neuroectoderm. At this stage, the anterior gut endoderm is coalescing at the embryonic midline in wild-type embryos. In *fau* mutants, gut endoderm is only present laterally (arrow). For purposes of orientation, the arrow is at the level of the midbrain–hindbrain boundary. (B) *fkd2* is expressed throughout the pancreas (arrowhead), liver (arrow), and posterior gut tube in wild-type embryos. *fau* mutants display variable abnormalities in the morphogenesis and amount of *fkd2*-positive tissue. These defects range from near absence of *fkd2*-positive endoderm (top right) to formation of two lateral gut tubes (bottom left) to the absence of gut looping (bottom right). (C) Morphogenesis of the pharyngeal pouches is disrupted in *fau* mutants. The mediolateral distance between the pouches is greater in *fau* mutants and the anterior pouches (arrow) are disorganized. Also, midline *nkx2.3*-expressing cells (arrowhead) appear to be absent in *fau* mutants. (D) In *fau*^{s26} mutants, the anterior domain of *nkx2.3* expression (arrow) is absent.

tion fragment length polymorphism (RFLP) within the *gata5* gene demonstrated that the mutant *gata5* allele segregated with the *fau*^{tm236a} phenotype in >300 meioses.

Sequencing of the *fau*^{tm236a} allele of *gata5* revealed a G-to-A substitution at the first base of the third intron (Fig. 4A). This mutation disrupts the highly conserved GT dinucleotide necessary for recognition of the splice site by the U1 snRNP (Krawczak et al. 1992). Sequencing of *gata5* cDNAs from *fau* mutants demonstrated that the *fau*^{tm236a} mutation leads to mis-splicing at flanking cryptic splice sites (Fig. 4B). The most frequently isolated cDNA results from the use of a downstream cryptic splice site and contains a 31-bp insertion predicted to

create a translational frameshift (Fig. 4C,D). A less frequently isolated cDNA is produced by use of a cryptic splice site 30 bp upstream of the inactivated site (Fig. 4B). This mutant splice variant is predicted to produce a 10-amino-acid deletion in the second zinc finger (Fig. 4C,D).

Structure/function studies have revealed that murine Gata5 contains two amino-terminal transcriptional activation domains, as well as a basic domain required for nuclear localization (Morrissey et al. 1997b). Murine Gata4 also contains a carboxy-terminal transcriptional activation domain required for synergistic activity with Nkx2.5 (Durocher et al. 1997; Lee et al. 1998; Sepulveda et al. 1998). As Gata5 shows a similar synergism with



Nkx2.5 (Durocher et al. 1997), Gata5 may also contain a carboxy-terminal activation domain. Sequence comparison of murine Gata5 and zebrafish Gata5 suggests that the two proteins are similar in structure (Rodaway et al. 1999). The more prevalent 31-bp insertion caused by the *fau^{tm236a}* mutation disrupts the putative carboxy-terminal activation domain and the basic nuclear localization domain of zebrafish Gata5 (Fig. 4C,D). The rarer splicing event deletes a 30-bp region that encodes part of the carboxy-terminal zinc finger required for DNA binding and for interaction with the transcription factors NF-AT3 and Nkx2.5 (Durocher et al. 1997; Lee et al. 1998; Molkentin et al. 1998). Additionally, the carboxy-terminal zinc finger has also been implicated in mediating heterotypic interactions between Gata factors (Charron et al. 1999). Therefore, we suspect that both mutant forms of Gata5 retain little, if any, activity. However, wild-type message can be detected in *fau^{tm236a}* mutants (see Materials and Methods), indicating that the splicing machinery is still able to recognize the mutant splice site, albeit at a much reduced frequency. Taken together, these data indicate that *fau^{tm236a}* is a mutation in the *gata5* gene that substantially reduces, but does not eliminate, the production of wild-type Gata5.

Examination of the *fau^{s26}* *gata5* cDNA sequence did not identify a mutation. As *gata5* expression is dramatically diminished in *fau^{s26}* mutants (see below), we hypothesize that the *fau^{s26}* mutation disrupts the function of the *gata5* promoter. We sequenced 800 bp upstream of the transcriptional start site and have identified a putative core promoter without successfully identifying the *fau^{s26}* lesion (data not shown). Therefore, we speculate that the *fau^{s26}* mutation disrupts a distal enhancer of the *gata5* gene that reduces *gata5* transcription to very low levels.

gata5 expression in wild-type and *fau* mutant embryos

In zebrafish, *gata5* is first expressed at the dome stage (late blastula) in the most marginal cells (Fig. 5A, arrowhead), a population that includes the endodermal progenitors and some mesodermal progenitors (Warga and Nüsslein-Volhard 1999), and in the yolk syncytial layer (YSL), an extraembryonic tissue underlying the blastoderm that may be equivalent to the mouse visceral endoderm (Fig. 5A, arrow). During gastrulation, *gata5*-expressing cells in the hypoblast adopt one of two different morphologies. Predominantly dorsally, *gata5*-expressing cells adopt the large, flat morphology characteristic of endoderm (Fig. 5C, arrowhead) (Warga and Nüsslein-Volhard 1999). Further evidence that these large *gata5*-expressing cells are endodermal progenitors comes from the very similar distribution of these cells and the *axial*- and *sox17*-expressing endodermal progenitors (Alexander and Stainier 1999). In a more ventral domain reflecting the regional propensity to become heart (Stainier et al. 1993), other *gata5*-expressing cells adopt a rounder morphology characteristic of mesodermal progenitors (Fig. 5C, arrow). After completion of gastrulation, *gata5* expression persists in the endo-




Figure 5. *gata5* expression in zebrafish embryos at dome (4.3 hpf; A), germ ring (5.7 hpf; B), 90% epiboly (9 hpf; C), bud (10 hpf; D), 4-somite (11.3 hpf; E), 28 hpf (F,H), and 80% epiboly (8.3 hpf; G) stages. A,B,C, and G show lateral views with dorsal to the right. D,E,F, and H show dorsal views with anterior to the top. (A) At dome stage, *gata5* is expressed in the yolk syncytial layer (arrow) and the most marginal tiers of the blastoderm (arrowhead). (B) During early gastrulation, *gata5* expression is maintained in the marginal zone (bracket) and yolk syncytial layer (arrow). (C) Late in gastrulation, *gata5* is expressed by cells of two different morphologies. Predominantly dorsally, endodermal cells displaying a characteristically large and flat morphology (arrowhead) express *gata5*. Ventrally, hypoblast cells with a rounder, more mesodermal appearance (arrow) also express *gata5*. (D) After the completion of gastrulation, *gata5* is expressed in the endoderm (stippled staining) and ALPM (arrow). (E) During early somitogenesis, both the endoderm and lateral plate mesoderm (arrow) continue to express *gata5*. (F) At 28 hpf, *gata5* is expressed most strongly in the endocardium (arrow) and weakly in the surrounding noncardiogenic ALPM. (G) *fau^{s26}* mutants display markedly reduced *gata5* expression. At 80% epiboly, *gata5* mRNA is undetectable in the YSL and observed only faintly (arrow) in the blastoderm of *fau^{s26}* mutants. (H) *gata5* expression is undetectable in *fau^{s26}* mutants at 28 hpf.

derm and ALPM (Fig. 5D). ALPM expression of *gata5* is maintained throughout somitogenesis (10–24 hpf) whereas the level of endodermal expression diminishes (Fig. 5E). Later, from 20–22 hpf, cardiac expression of *gata5* becomes predominantly endocardial (Fig. 5F, arrow), as has been described for murine *gata5* (Morrissey et al. 1997a).

Expression of *gata5* in *fau^{tm236a}* mutants and wild-type siblings is indistinguishable until morphogenetic defects become apparent. In contrast, *fau^{s26}* mutants display markedly decreased *gata5* expression throughout development and in both the YSL and embryonic tissues (Fig. 5G,H). Very low levels of *gata5* expression can be detected in *fau^{s26}* mutants by in situ hybridization (Fig. 5G, arrow) and RT-PCR (data not shown), indicating that although expression is strikingly diminished, the *fau^{s26}* mutation does not completely eliminate the production of *gata5* message.

Overexpression of wild-type *gata5* rescues heart fusion in *fau^{tm236a}* mutants

To confirm that the *fau* mutant phenotype is due to the absence of Gata5 activity, we injected mRNA encoding wild-type Gata5 into *fau^{tm236a}* mutants at the 1–4 cell stage. Although this approach leads to widespread misexpression, injection of 20 pg of *gata5* mRNA restores cardiac fusion in *fau^{tm236a}* mutants (Fig. 6A; Table 1), confirming that the *fau* phenotype is due to a loss-of-function mutation in *gata5*. *gata5*-injected *fau* mutants are not rescued to viability as *gata5* misexpression causes variable abnormalities in morphogenesis, including anterior truncations and compression of the anterior–posterior axis. Injection of *tm236a-1*, a *gata5* mRNA containing the same insertion and frameshift as most commonly caused by the *fau^{tm236a}* mutation, does not rescue cardiac fusion (Table 1). Moreover, *tm236a-1* overexpression does not disrupt morphogenesis, indicating that it has little if any biological activity.

To test whether the defects in organogenesis exhibited by *fau* mutants are due to reduced Gata5 activity in the extraembryonic YSL, we injected *gata5* mRNA directly into the YSL. Overexpression of *gata5* mRNA in the YSL does not restore cardiac fusion in *fau^{tm236a}* mutants (14 cardia bifida mutants observed in a clutch of 60 YSL-injected embryos), suggesting that *gata5* is required embryonically to promote heart fusion.

Ectopic expression of *gata5* leads to the ectopic expression of myocardial genes

Injection of wild-type embryos with 50 pg of *gata5* mRNA expands the ALPM expression of *cmlc1*, *cmlc2*, *vmhc*, and the MF20 and S46 myosin heavy chain proteins and leads to their ectopic expression (Fig. 6B,C; Table 2; data not shown). Occasionally, ectopic domains of myocardial gene expression both in the head and the tail were observed to contract rhythmically (Table 2). Control embryos injected with *gfp* mRNA did not manifest either expansion of myocardial gene expression or ectopic regions of beating tissue.

Because *nkx2.5* is downregulated in *fau* mutants, we investigated whether *gata5* overexpression was sufficient to induce the expression of zebrafish *nkx2.5* or other myocardial regulatory genes. Notably, we found that injection of 50 pg of *gata5* mRNA expands the do-




Figure 6. Overexpression of *gata5* rescues cardiac fusion in *fau^{tm236a}* mutants and leads to ectopic expression of myocardial genes. Expression of *cmlc1* (A,C), *cmlc2* (B), *nkx2.5* (D), *gata4* (E), and *gata6* (F) in wild-type embryos and *fau* mutant siblings at 36 hpf (A), 25 hpf (B,C), 5-somite stage (11.7 hpf; D), and 12-somite stage (14.5 hpf; E,F). Views are anterior [A], lateral [B,C,D (bottom right)], and dorsal [D (top, and bottom left), E,F]. (A) Injection of 20 pg of *tm236a-1* mRNA has no demonstrable effect on development (top left) and does not restore cardiac fusion in *fau^{tm236a}* mutants (top right, arrows). In contrast, injection of an equivalent amount of wild-type *gata5* mRNA causes variable defects in head and body morphology (bottom left) and can restore cardiac fusion in *fau^{tm236a}* mutants (bottom right, arrowhead). (B) Injection of a higher dose of *gata5* mRNA (50 pg) can cause expanded expression of *cmlc2* in a domain contiguous to the ALPM. (C) *gata5* overexpression can also cause ectopic expression of *cmlc1*. Here, *gata5* overexpression has led to ectopic *cmlc1* expression in cells overlying the ventral side of the yolk (arrow). A heart tube has also formed in the expected location (arrowhead). (D) Injection of 50 pg of *gata5* mRNA can increase the number of cells expressing *nkx2.5* (top right) and cause ectopic *nkx2.5* expression (bottom). The bottom panels show dorsal and lateral views of the same embryo, which exhibits a dorsal midline region of ectopic *nkx2.5* expression (arrows). β -Galactosidase staining produces a turquoise color marking the distribution of coinjected *lacZ* mRNA. *gata5* overexpression can also lead to increased expression of *gata4* (E) and *gata6* (F).

main of *nkx2.5*, *gata4*, and *gata6* expression within the ALPM (Fig. 6D–F; Table 2) and causes them to be ectopically expressed (Fig. 6D; Table 2). Coinjection of *lacZ* and *gata5* mRNA allowed us to correlate the embryonic distribution of the exogenous RNA with the increased expression of *nkx2.5*. In all cases but one ($n = 131$), regions of expanded or ectopic *nkx2.5* expression overlapped with the distribution of exogenous *gata5* mRNA as indicated by β -galactosidase activity (Fig. 6D; Table 2),

Table 1. Overexpression of wild-type *gata5* restores cardiac fusion in *fau^{tm236a}* mutants

Injected mRNA	Number of <i>fau^{tm236a}</i> mutants (by genotype)	Number with cardia bifida
<i>gata5</i>	39	9 (23%)
<i>tm236a-1</i>	25	19 (76%)
Uninjected	35	23 (70%)

The observed value of the test $\chi^2(1)$, 17.33, indicates that the increased rate of cardiac fusion following overexpression of *gata5* in *fau^{tm236a}* mutants is statistically significant.

suggesting that Gata5 autonomously induces the expression of myocardial genes.

Surprisingly, the ectopic expression of these genes does not appear to be limited to the mesodermal germ layer. Ectopic expression of *nkx2.5*, *gata4*, and *gata6* was observed in dorsal regions of the embryo normally fated to form neuroectoderm (Fig. 6D; data not shown). Neuroectodermal expression of *nkx2.5* and *gata6* is never observed during normal zebrafish embryogenesis.

Human *gata5* maps to 20q13.2–q13.3

Mutations in human *NKX2-5* can cause congenital cardiac septation defects (Schott et al. 1998). As Gata4 and Gata5 physically interact and synergize with *Nkx2.5*

Table 2. *gata5* overexpression causes ectopic expression of myocardial genes and beating tissue

Injected mRNA	Riboprobe	Expanded expression	Ectopic expression
<i>gata5</i>	<i>cmlc1</i>	32/95	10/95
<i>gata5</i>	<i>cmlc2</i>	20/64	7/64
<i>gata5</i>	<i>vmhc</i>	13/22	4/22
<i>gata5</i>	<i>nkx2.5</i>	11/18	4/18
<i>gata5</i>	<i>nkx2.5</i>	0/6	0/6
<i>gata5</i> and <i>lacZ</i>	<i>nkx2.5</i>	84/237 ^a	47/237 ^a
<i>gfp</i> and <i>lacZ</i>	<i>nkx2.5</i>	0/36	0/36
<i>gata5</i>	<i>gata4</i>	9/16	6/16
<i>gfp</i>	<i>gata4</i>	0/6	0/6
<i>gata5</i>	<i>gata6</i>	14/22	10/22
<i>gfp</i>	<i>gata6</i>	0/5	0/5
Ectopic beating tissue			
<i>gata5</i>		4/52	
<i>gfp</i>		0/7	

Embryos were scored as displaying expanded expression if expression was noted in cells adjacent to the normal domain [see Fig. 6B,D (upper right), E,F for examples]. Embryos were scored as displaying ectopic expression or ectopic beating tissue if tissue was noted dorsally or posteriorly within the embryo in a domain discontinuous with the ALPM or heart [see Fig. 6C,D (bottom) for examples].

^aOf the *gata5*-overexpressing embryos exhibiting expanded or ectopic *nkx2.5* expression, 84/84 and 46/47 of the embryos, respectively, displayed increased expression within a domain of β -galactosidase activity.

(Durocher et al. 1997; Lee et al. 1998; Sepulveda et al. 1998), and as zebrafish Gata5 is essential for the correct expression of *nkx2.5*, it is possible that mutations in *gata* genes also cause human congenital heart defects. In support of this hypothesis, *gata4* haploinsufficiency has been associated with congenital heart malformations (Pehlivan et al. 1999). Human *gata4* and *gata6* map to chromosome 8p23 (Pehlivan et al. 1999) and 18q11.1–q11.2 (Suzuki et al. 1996), respectively. Using the Stanford G3 radiation hybrid panel, we mapped human *gata5* to chromosome 20q13.2–q13.3. The study of appropriate pedigrees should determine whether *gata5* also plays an important role in human congenital heart disease.

Discussion

gata5 is required for cardiac morphogenesis and differentiation, as well as the production of normal amounts of myocardial tissue

In this study we establish that the *fau* locus encodes zebrafish Gata5, an evolutionarily conserved zinc finger transcription factor, and that mutations in *gata5* profoundly disrupt cardiac development. Most conspicuously, diminished Gata5 activity prevents the precardiac mesoderm from migrating to and fusing at the embryonic midline, resulting in cardia bifida.

Reduced Gata5 activity also disrupts the expression of several genes in the ALPM, including *nkx2.5*. The defect in *nkx2.5* expression is observable early in somitogenesis indicating that Gata5 is required very early for proper myocardial differentiation. As previous studies have demonstrated that two murine *nkx2.5* regulatory elements contain essential Gata-binding sites (Searcy et al. 1998; Lien et al. 1999), Gata5 may be a direct endogenous regulator of *nkx2.5*. Consistent with this hypothesis, *gata5* is expressed during blastula stages in a domain that includes the myocardial precursors, whereas we first observe *nkx2.5* expression in a subset of *gata5*-expressing cells only after the completion of gastrulation (data not shown). Similar relative timing of expression of these genes is seen during chick development (T. Evans, pers. comm.).

Gata5 is capable of activating transcription from a wide range of myocardial promoters in vitro (Charron and Nemer 1999). Our results suggest that the same is true in vivo; *gata5* mutants display marked defects in the expression of many genes encoding components of the myocardial sarcomere (e.g., *cmlc1*, *cmlc2*, *vmhc*, *cardiac troponin T*, *tropomyosin*). However, it is not clear at this point which of these genes are direct targets of Gata5 and which require intermediate Gata5-dependent regulators. In fact, there may be no clear division between these cases as Gata5 may act both directly and indirectly on a single promoter. For example, Gata5 may participate in the induction of *nkx2.5* and also bind cooperatively with *Nkx2.5* to regulatory elements of a wide range of myocardial genes. It is also interesting to note the different requirements various sarcomeric protein genes have for Gata5. For example, although the expression patterns of

cmlc1 and *cmlc2* are indistinguishable in wild-type embryos, *cmlc1* expression is more severely reduced than *cmlc2* expression in *fau* mutants. Perhaps these differences reflect different affinities of Gata5 for the respective *cis* regulatory elements.

In addition to exhibiting defects in myocardial differentiation, the hearts of *fau* mutants at 33 hpf are very small in comparison to those of wild-type siblings and other cardia bifida mutants such as *cas*. Similarly, the number of developing myocardial precursors is reduced in *fau* mutants throughout earlier segmentation stages. This deficit does not seem to reflect problems in ALPM specification or maintenance as *gata6* expression demonstrates that the ALPM is present and morphologically normal in *gata5* mutants. Instead, Gata5 appears to play a role in the induction, differentiation, and/or proliferation of the myocardial precursors within the ALPM.

Differentiation of the myocardium into atrial and ventricular components is also defective in *fau* mutants. *fau* myocardia preferentially form atrial tissue, indicating that *gata5* is required for ventricular induction or differentiation and that the two chambers of the heart have different requirements for *gata5*. However, zebrafish *gata5* does not appear to be differentially expressed in the two chambers of the heart. Although we have not examined the distribution of Gata5 protein, the expression pattern of the *gata5* gene suggests that it is not directly involved in patterning the cardiac chambers. Instead, the defect in ventricular development may be secondary to the reduced amount of myocardial tissue produced in *gata5* mutants if embryos must elaborate a threshold amount of myocardial tissue to sustain ventricular induction or differentiation. Alternatively, Gata5 may act with an unidentified ventricle-specific factor to promote ventricular development.

Taken together, these results demonstrate that Gata5 has a critical role in multiple aspects of zebrafish heart development including the expression of *nkx2.5* and other myocardial genes, the production of a normal amount of myocardial precursors, the morphogenesis of both the myocardial and endocardial components of the heart, and the differentiation of the ventricular chamber.

gata5 is required for endoderm morphogenesis

Previous expression studies have suggested a role for Gata factors in the early development of the embryonic endoderm (Laverriere et al. 1994; Morrissey et al. 1997a; Bossard and Zaret 1998). Here, we have confirmed that zebrafish *gata5* is essential for the formation of normal amounts and morphogenesis of the endoderm. Although gut and pharyngeal endoderm lineages separate early during zebrafish development and have distinct genetic requirements (Alexander 1999), *gata5* is necessary for the morphogenesis of both tissues. Interestingly, Gata5 is required for the movement toward the midline of both the heart primordia and the gut and pharyngeal endoderm, suggesting a causative link between these defects or a common role of Gata5 in promoting the medial migration of several tissues.

gata5 overexpression causes the ectopic expression of myocardial genes

We show that overexpression of zebrafish *gata5* leads to the expanded or ectopic expression of *cmlc1*, *cmlc2*, *vmhc*, *gata4*, *gata6*, *nkx2.5*, and the MF20 and S46 myosin heavy chain proteins, indicating that zebrafish Gata5 can act as a positive regulator of many myocardial genes, even in noncardiogenic tissues. This is the first demonstration of the ability of a Gata factor to enhance cardiogenesis in the context of a developing embryo. In contrast, overexpression of *gata4* and *gata5* in *Xenopus* does not affect the spatial pattern of myocardial gene expression or heart morphogenesis (Jiang and Evans 1996). The apparently different capacities of *Xenopus* and zebrafish *gata5* may reflect species-dependent differences in the plasticity of cell fate or differences in the activities of the two homologs. In fact, zebrafish Gata5 function may be more similar to that of murine Gata4, which can also potentiate cardiogenesis, at least in the P19 cell line (Grepin et al. 1997).

The ability of Gata5 to induce ectopic myocardial gene expression is not entirely unexpected, given that Gata5 regulates *nkx2.5* expression and overexpression of *nkx2.5* can induce some ectopic myocardial gene expression (Chen and Fishman 1996). It will be interesting to determine whether Nkx2.5 is an obligatory effector of Gata5 function in myocardial cells. Another intriguing issue concerns whether Gata5, when forcibly expressed in noncardiogenic tissues, simply activates the transcription of a subset of myocardial genes or whether Gata5 can reprogram an otherwise noncardiac cell to become completely myocardial in character. The unique ability of *gata5* overexpression to produce ectopic beating tissue suggests that, at least in some contexts, these cells exhibit myocardial behavior. Therefore, both phenotypic analysis of *fau* mutants and gain-of-function experiments indicate that Gata5 plays an essential role in the induction, differentiation, and/or proliferation of the myocardial precursors.

Although the normal expression of *gata5* in *fau*^{tm236a} mutants suggests that Gata5 does not regulate its own expression, *fau* mutants exhibit modestly decreased expression of *gata4*, and overexpression of *gata5* is able to induce the ectopic transcription of *gata4*. Therefore, Gata5 appears to positively regulate *gata4*, through which Gata5 may mediate some of its effects on cardiac differentiation. Although no defects in *gata6* expression were noted in *fau* mutants, *gata5* overexpression can cause ectopic *gata6* expression, suggesting that Gata5 may have a redundant role in regulating *gata6*. Perhaps embryos mutant for both *gata4* and *gata5* will reveal a nonessential role for Gata5 in *gata6* gene regulation. Previous work on mouse *gata4* and *gata6* mutants has demonstrated that Gata4 negatively regulates *gata6* in the ALPM whereas Gata6 positively regulates *gata4* in extraembryonic tissues (Kuo et al. 1997; Molkentin et al. 1997; Morrissey et al. 1998). Thus, complex interregulation by Gata4, Gata5, and Gata6 may have an important

role in controlling the early events of cardiac and endodermal development.

gata5 may act both tissue autonomously and nonautonomously in heart development

Previous studies in both mouse and zebrafish have demonstrated that cardia bifida can be secondary to defects in endoderm development (Narita et al. 1997; Peyrieras et al. 1998). This work leads us to hypothesize that the cardia bifida in *fau* mutants is also secondary to their severe endodermal defects. Moreover, the presence of endodermal defects in the cardia bifida mutants *fau*, *oep*, and *cas* (Schier et al. 1997; Alexander et al. 1999) suggests that zebrafish cardia bifida may be a general consequence of defects in endodermal morphogenesis or differentiation. Conclusive experimental tests of the autonomy of the morphogenetic defects in *fau* mutants will require analyses of embryos in which interacting tissues are of distinct genotypes, the production of which is contingent on techniques not yet developed in zebrafish.

Although the absence of cardiac fusion in *fau* mutants may be secondary to endoderm defects, other cardiac abnormalities may be due to reduced Gata5 activity in the ALPM. Despite the complete and early absence of gut endoderm in *cas* mutants, *cas* myocardia exhibit no defect in gene expression (Alexander et al. 1999), are larger than *fau* myocardia, and contain a normal ratio of atrial to ventricular tissue. Thus, *fau* mutants display defects in gene expression, cardiac growth, and ventricular development beyond what is attributable to the loss of endoderm, implicating ALPM *gata5* expression in these roles. The observation that *gata5* overexpression autonomously promotes the ectopic expression of *nkx2.5* further suggests that Gata5 acts within cells of the ALPM to regulate myocardial differentiation. We therefore propose that the defects in *fau* cardiac morphogenesis are secondary to the reduction of Gata5 activity in the endoderm, whereas the additional defects in cardiac growth and differentiation are due to the reduction of Gata5 function in the ALPM.

Zebrafish Gata5 and murine Gata4 may function similarly during organogenesis

Although mouse *gata5* null mutants have not been extensively described, they are viable (Morrissey et al. 1998), suggesting that some of the functions performed by zebrafish Gata5 may be fulfilled by Gata4 or Gata6 in the mouse. Unfortunately, mouse embryos deficient for Gata4 (Kuo et al. 1997; Molkentin et al. 1997) or Gata6 (Morrissey et al. 1998; Koutsourakis et al. 1999) arrest at embryonic day (E) 7.0–9.5 and E 5.5–7.5, respectively, impeding the analysis of later developmental events such as cardiac chamber formation and the development of endoderm-derived organs. However, *gata4* null mice and *gata5* mutant zebrafish both display cardia bifida, prompting a comparison. *gata4* null mice do not undergo

ventral closure, which has led to the idea that cardia bifida in *gata4* null mice is a direct consequence of the defect in ventral folding. Our study, indicating that zebrafish cardia bifida can result from *gata*-dependent defects in the migration of the cardiac primordia to the embryonic midline, suggests an alternative interpretation. In zebrafish, the cardiac precursors migrate on the yolk syncytial layer to reach the embryonic midline (Stainier et al. 1993). Perhaps murine precardiac mesoderm reaches the embryonic midline similarly, by actively migrating on the underlying visceral endoderm and not through ventral folding of the body wall. In support of this hypothesis, in the mouse the midline cardiogenic plate appears to form before the beginning of ventral folding (Plate 8d in Kaufman 1992; Saga et al. 1999). Therefore, cardia bifida in *gata4* null mice may stem from a defect in the migration of the precardiac mesoderm to the embryonic midline in much the same way as cardia bifida arises in mice lacking MesP1 (Saga et al. 1999). We propose that mouse *gata4* is required for two early and distinct morphogenetic processes: (1) the initial migration of the bilateral cardiac precursors to the embryonic midline where they form the cardiogenic plate, and (2) the subsequent ventral folding required for foregut and ventral body-wall morphogenesis. Determining whether cardiac fusion is truly independent of ventral closure will require the identification of mouse mutations that inhibit ventral closure without affecting cardiac fusion.

In summary, we have shown that *gata5* mutants show reduced expression of *nkx2.5* and that *gata5* overexpression leads to ectopic *nkx2.5* expression. These data together with the demonstrated presence of Gata-binding sites in *nkx2.5* regulatory elements indicate that *gata5* functions upstream of *nkx2.5* to initiate myocardial differentiation. Further studies will be required to identify additional regulators of heart and endoderm development that function downstream of or in combination with Gata5.

Materials and methods

Zebrafish strains and maintenance

Zebrafish were maintained and staged as described (Westerfield 1995).

Whole-mount in situ hybridization and immunohistochemistry

We performed in situ hybridization as described (Alexander et al. 1998); in certain cases, embryos older than 28 hpf were incubated in 0.003% phenylthiourea to inhibit pigmentation. The *gata4* (GenBank accession no. AF191577) and *gata6* (GenBank accession no. AF191578; previously called *gata5* in Pack et al. 1996) genes were isolated in Leonard Zon's laboratory (Harvard Medical School, Boston, MA; unpubl.). Full-length cDNA was used as the template for the *gata6* riboprobe, but the *gata4* riboprobe was generated from a truncated cDNA comprised of the first 954 bp. The *gata5* riboprobe is described below. The *cmlc1* gene (GenBank accession no. AF119163) was isolated as

described (Yelon et al. 1999). As *fau* mutants younger than 24 hpf could not be phenotypically distinguished from wild-type siblings, earlier stage embryos were genotyped post in situ hybridization to confirm that any defects in gene expression segregated with the *fau* mutation. Briefly, this entailed serial rehydration with PBS + 0.1% Tween, proteinase K digestion at 55°C overnight, and PCR genotyping with a tightly linked marker.

Immunohistochemistry using the antibodies MF20 and S46 (generous gift of F. Stockdale, Stanford University, Palo Alto, CA) was performed as described (Alexander et al. 1998). MF20 was obtained from the Developmental Studies Hybridoma Bank, maintained by the Department of Biological Studies, University of Iowa, Ames under contract NO1-HD-2-3144 from the National Institute of Child Health and Development (NICHD).

β -Galactosidase staining was performed as described (Rodaway et al. 1999).

Genetic mapping

The mapping strain was constructed by crossing a *fau* Tü male heterozygote to a WIK female. Genomic DNA extraction from zebrafish embryos was performed as described (Westerfield 1995), except that proteinase K digestion was done for 2–4 hr. By using only those embryos recombinant at simple sequence length polymorphisms flanking *fau* in a search for AFLPs, we successfully restricted our identified markers to those that mapped within a useful genetic distance of *fau*. Otherwise, AFLP selection was performed as described (Vos et al. 1995). An AFLP marker linked to the *fau* mutation (primers: 5'-AAGAGAGCGAGCACTGCTACTAC-3' and 5'-CAGCTTTC-CATATTCTGGC-3') and an amplicon from the *gata5* 3' UTR (primers: 5'-TGATCCGAGACCTCCAC-3' and 5'-CTCCC-GATCATTGGAAC-3') were placed on the Goodfellow zebrafish radiation hybrid panel (Research Genetics). These markers are separated by 14 cR (lod score 18.96). Human *gata5* was mapped using primers designed from the sequence of EST W00677 (primers: 5'-AGACCACCCACTGAGTCACC-3' and 5'-GGAGTCCCTTGCTGTACGTG-3') on the Stanford G3 radiation hybrid panel (Research Genetics). Human *gata5* is linked to marker SHGC-57473 with a lod score of 18.28.

PAC library screen and linkage analysis

A PAC library was screened by PCR as pools of clones using primers that amplify the *gata5* 3' UTR. Internal PAC sequencing was performed using the primer 5'-ACACTGTGGAG-GAGAAACGCA-3'. A single-strand conformational polymorphism identified at one end of a PAC clone containing *gata5* was linked to the *fau^{tm236a}* mutation in all of 272 gynogenetic haploid embryos examined. Additionally, a BsmFI RFLP at position 380 in the *gata5* gene was linked to *fau^{tm236a}* in all of 160 diploid embryos examined. The 400-bp region containing the RFLP was amplified by PCR (primers: 5'-GTACCAGCCAT-GCTCCAGAC-3' and 5'-CTCCTTCCCGGTAGAGTTCC-3'). The *fau^{s26}* mutation was linked to SSRP z4003 in all of 40 gynogenetic haploid embryos examined. Amplification of SSRPs flanking *gata5* from *fau^{s26}* genomic DNA failed to reveal evidence of a deletion.

Isolation of *gata5* cDNA

Total RNA was isolated using Trizol (Gibco BRL). Reverse transcription was performed using Superscript II (Gibco BRL). The *gata5* ORF was amplified by PCR (primers: 5'-GGACGTT-GACAAGGTTTTTACTAG-3' and 5'-GTCTGTCTACATTT-GTGTAAGTCC-3'). The 1330-bp product was cloned into

the pGEM-T vector (Promega) and sequenced using an ABI 377 DNA sequencer. We sequenced 20 independent clones of mutant-derived and 8 independent clones of wild-type-derived *gata5* cDNA (one of which was used as the template for the *gata5* riboprobe). Among the mutant-derived clones, the 31-bp insertion was represented by 17 clones, the 30-bp deletion was represented by 2 clones, and the wild-type cDNA was detected once. PCR analysis using a primer bridging the wild-type splice site (primer: 5'-GCTTCATGGGGTACCAAGG-3') confirmed that wild-type *gata5* message is detectable in *fau^{tm236a}* mutants.

Although the amount of *gata5* mRNA is profoundly diminished in *fau^{s26}* mutants, low levels were detectable by RT-PCR performed as above. Sequencing of the subcloned RT-PCR products indicated that *fau^{s26}* is not a mutation in the *gata5* coding sequence or UTRs.

gata5 gene structure

PCR of genomic amplicons and subsequent sequencing revealed that the first intron (723 bp) is present 3' of cDNA nucleotide 602, the second intron (1207 bp) is present 3' of cDNA nucleotide 771, the third intron (1.6 kb) is present 3' of cDNA nucleotide 901, and the fourth intron (343 bp) is present 3' of cDNA nucleotide 986. Additional internal PAC sequencing upstream of the transcriptional start site using the primer 5'-GGCTAGT-TAAAAACCTTGTCACG-3' revealed a putative promoter sequence containing a consensus TATA box and several potential Gata-binding elements. Sequencing of this region in *fau^{s26}* mutants failed to identify a mutation.

gata5 mRNA injection

Full-length, capped mRNA was generated from pCS2:*gata5*, pCS2:tm236-1:MT, and pGST:GFP using the SP6 mMESSAGE mMACHINE system (Ambion). pCS2:tm236-1:MT was created by subcloning the coding sequence of the *gata5* cDNA species most frequently isolated from *fau^{tm236a}* mutants into pCS2:MT. For embryonic overexpression, 20 or 50 pg of mRNA was microinjected at the 1–4 cell stage. For autonomy studies, 100 pg of *lacZ* mRNA was coinjected. To restrict overexpression to the YSL, mRNA was microinjected directly into the YSL at the 512-cell stage. Injected embryos were raised to 11–36 hpf for in situ hybridization or to 30 hpf for immunohistochemical analysis. After phenotypic analysis, embryos were genotyped as described above.

Acknowledgments

We dedicate this paper to the memory of Nigel Holder. We thank Sarah Howard, Ken Nakashima, Alex Navarro, and Elliot Perens for invaluable assistance; members of the Stainier laboratory, Brian Black, Todd Evans, Richard Harvey, Ira Herskowitz, and Patrick Tam for helpful discussions and/or comments; Joseph DeYoung and Travis Taylor for sequencing; Len Zon for providing *gata4* and *gata6*; and Frank Stockdale for generously providing the S46 antibody. J.F.R. and J.A. are members of the Medical Scientist Training Program of the National Institutes of Health. D.Y. is an Amgen fellow of the Life Sciences Research Foundation. This work was supported by the American Heart Association (J.A. and D.Y.R.S.), the Medical Research Council and Wellcome Trust (A.R., N.H., and R.P.), the Packard Foundation and the Life and Health Insurance Medical Research Fund (D.Y.R.S.).

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby

marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Alexander, J. 1999. "Gut feelings: Molecular and genetic analysis of endoderm development in zebrafish", Ph.D. Thesis. University of California, San Francisco, San Francisco, CA.
- Alexander, J. and D.Y.R. Stainier. 1999. A molecular pathway underlying endoderm formation in zebrafish. *Curr. Biol.* **9**: 1147–1157.
- Alexander, J., D.Y.R. Stainier, and D. Yelon. 1998. Screening mosaic F1 females for mutations affecting zebrafish heart induction and patterning. *Dev. Genet.* **22**: 288–299.
- Alexander, J., M. Rothenberg, G.L. Henry, and D.Y.R. Stainier. 1999. *casanova* plays an early and essential role in endoderm formation in zebrafish. *Dev. Biol.* **215**: 343–357.
- Bossard, P. and K.S. Zaret. 1998. GATA transcription factors as potentiators of gut endoderm differentiation. *Development* **125**: 4909–4917.
- Charron, F. and M. Nemer. 1999. GATA transcription factors and cardiac development. *Semin. Cell. Dev. Biol.* **10**: 85–91.
- Charron, F., P. Paradis, O. Bronchain, G. Nemer, and M. Nemer. 1999. Cooperative interaction between GATA-4 and GATA-6 regulates myocardial gene expression. *Mol. Cell. Biol.* **19**: 4355–4365.
- Chen, J.-N. and M.C. Fishman. 1996. Zebrafish *tinman* homolog demarcates the heart field and initiates myocardial differentiation. *Development* **122**: 3809–3816.
- Chen, J.-N., P. Haffter, J. Odenthal, E. Vogelsang, M. Brand, F.J. van Eeden, M. Furutani-Seiki, M. Granato, M. Hamerschmidt, C.P. Heisenberg et al. 1996. Mutations affecting the cardiovascular system and other internal organs in zebrafish. *Development* **123**: 293–302.
- Durocher, D., F. Charron, R. Warren, R.J. Schwartz, and M. Nemer. 1997. The cardiac transcription factors Nkx2-5 and GATA-4 are mutual cofactors. *EMBO J.* **16**: 5687–5696.
- Evans, S.M. 1999. Vertebrate tinman homologues and cardiac differentiation. *Semin. Cell. Dev. Biol.* **10**: 73–83.
- Evans, T. 1997. Regulation of cardiac gene expression by GATA-4/5/6. *Trends Cardiovasc. Med.* **7**: 75–83.
- Fukushige, T., M.G. Hawkins, and J.D. McGhee. 1998. The GATA-factor *elt-2* is essential for formation of the *Caenorhabditis elegans* intestine. *Dev. Biol.* **198**: 286–302.
- Gannon, M. and D. Bader. 1995. Initiation of cardiac differentiation occurs in the absence of anterior endoderm. *Development* **121**: 2439–2450.
- Gao, X., T. Sedgwick, Y.B. Shi, and T. Evans. 1998. Distinct functions are implicated for the GATA-4, -5, and -6 transcription factors in the regulation of intestine epithelial cell differentiation. *Mol. Cell. Biol.* **18**: 2901–2911.
- Gove, C., M. Walmsley, S. Nijjar, D. Bertwistle, M. Guille, G. Partington, A. Bomford, and R. Patient. 1997. Over-expression of GATA-6 in *Xenopus* embryos blocks differentiation of heart precursors [published erratum appears in *EMBO J.* 1997 **16**: 1806–1807]. *EMBO J.* **16**: 355–368.
- Grepin, C., G. Nemer, and M. Nemer. 1997. Enhanced cardiogenesis in embryonic stem cells overexpressing the GATA-4 transcription factor. *Development* **124**: 2387–2395.
- Jacobson, A.G. and A.K. Sater. 1988. Features of embryonic induction. *Development* **104**: 341–359.
- Jiang, Y. and T. Evans. 1996. The *Xenopus* GATA-4/5/6 genes are associated with cardiac specification and can regulate cardiac-specific transcription during embryogenesis. *Dev. Biol.* **174**: 258–270.
- Jiang, Y., S. Tarzami, J.B. Burch, and T. Evans. 1998. Common role for each of the cGATA-4/5/6 genes in the regulation of cardiac morphogenesis. *Dev. Genet.* **22**: 263–277.
- Kaufman, M.H. 1992. *The atlas of mouse development*. Academic Press, London, UK.
- Koutsourakis, M., A. Langeveld, R. Patient, R. Beddington, and F. Grosfeld. 1999. The transcription factor GATA6 is essential for early extraembryonic development. *Development* **126**: 723–732.
- Krawczak, M., J. Reiss, and D.N. Cooper. 1992. The mutational spectrum of single base-pair substitutions in mRNA splice junctions of human genes: Causes and consequences. *Hum. Genet.* **90**: 41–54.
- Kuo, C.T., E.E. Morrisey, R. Anandappa, K. Sigrist, M.M. Lu, M.S. Parmacek, C. Soudais, and J.M. Leiden. 1997. GATA4 transcription factor is required for ventral morphogenesis and heart tube formation. *Genes & Dev.* **11**: 1048–1060.
- Kwok, C., R.M. Korn, M.E. Davis, D.W. Burt, R. Critcher, L. McCarthy, B.H. Paw, L.I. Zon, P.N. Goodfellow, and K. Schmitt. 1998. Characterization of whole genome radiation hybrid mapping resources for non-mammalian vertebrates. *Nucleic Acids Res.* **26**: 3562–3566.
- Laverriere, A.C., C. MacNeill, C. Mueller, R.E. Poelmann, J.B. Burch, and T. Evans. 1994. GATA-4/5/6, a subfamily of three transcription factors transcribed in developing heart and gut. *J. Biol. Chem.* **269**: 23177–23184.
- Lee, Y., T. Shioi, H. Kasahara, S.M. Jobe, R.J. Wiese, B.E. Markham, and S. Izumo. 1998. The cardiac tissue-restricted homeobox protein Csx/Nkx2.5 physically associates with the zinc finger protein GATA4 and cooperatively activates *atrial natriuretic factor* gene expression. *Mol. Cell. Biol.* **18**: 3120–3129.
- Lien, C.L., C. Wu, B. Mercer, R. Webb, J.A. Richardson, and E.N. Olson. 1999. Control of early cardiac-specific transcription of *Nkx2-5* by a GATA-dependent enhancer. *Development* **126**: 75–84.
- Lyons, M.S., B. Bell, D.Y.R. Stainier, and K.G. Peters. 1998. Isolation of the zebrafish homologues for the tie-1 and tie-2 endothelium-specific receptor tyrosine kinases. *Dev. Dyn.* **212**: 133–140.
- Molkentin, J.D., Q. Lin, S.A. Duncan, and E.N. Olson. 1997. Requirement of the transcription factor GATA4 for heart tube formation and ventral morphogenesis. *Genes & Dev.* **11**: 1061–1072.
- Molkentin, J.D., J.R. Lu, C.L. Antos, B. Markham, J. Richardson, J. Robbins, S.R. Grant, and E.N. Olson. 1998. A calcineurin-dependent transcriptional pathway for cardiac hypertrophy. *Cell* **93**: 215–228.
- Morrissey, E.E., H.S. Ip, Z. Tang, M.M. Lu, and M.S. Parmacek. 1997a. GATA-5: A transcriptional activator expressed in a novel temporally and spatially-restricted pattern during embryonic development. *Dev. Biol.* **183**: 21–36.
- Morrissey, E.E., H.S. Ip, Z. Tang, and M.S. Parmacek. 1997b. GATA-4 activates transcription via two novel domains that are conserved within the GATA-4/5/6 subfamily. *J. Biol. Chem.* **272**: 8515–8524.
- Morrissey, E.E., Z. Tang, K. Sigrist, M.M. Lu, F. Jiang, H.S. Ip, and M.S. Parmacek. 1998. GATA6 regulates HNF4 and is required for differentiation of visceral endoderm in the mouse embryo. *Genes & Dev.* **12**: 3579–3590.
- Narita, N., M. Bielinska, and D.B. Wilson. 1997. Wild-type endoderm abrogates the ventral developmental defects associated with GATA-4 deficiency in the mouse. *Dev. Biol.* **189**: 270–274.
- Nascone, N. and M. Mercola. 1995. An inductive role for the

- endoderm in *Xenopus* cardiogenesis. *Development* **121**: 515–523.
- Odenthal, J. and C. Nüsslein-Volhard. 1998. fork head domain genes in zebrafish. *Dev. Genes Evol.* **208**: 245–258.
- Orkin, S.H. and L.I. Zon. 1997. Genetics of erythropoiesis: Induced mutations in mice and zebrafish. *Annu. Rev. Genet.* **31**: 33–60.
- Pack, M., L. Solnica-Krezel, J. Malicki, S.C. Neuhauss, A.F. Schier, D.L. Stemple, W. Driever, and M.C. Fishman. 1996. Mutations affecting development of zebrafish digestive organs. *Development* **123**: 321–328.
- Pehlivan, T., B.R. Pober, M. Brueckner, S. Garrett, R. Slauch, R. Van Rheeden, D.B. Wilson, M.S. Watson, and A.V. Hing. 1999. GATA4 haploinsufficiency in patients with interstitial deletion of chromosome region 8p23.1 and congenital heart disease. *Am. J. Med. Genet.* **83**: 201–206.
- Peyrieras, N., U. Strahle, and F. Rosa. 1998. Conversion of zebrafish blastomeres to an endodermal fate by TGF β -related signaling. *Curr. Biol.* **8**: 783–786.
- Rehorn, K.P., H. Thelen, A.M. Michelson, and R. Reuter. 1996. A molecular aspect of hematopoiesis and endoderm development common to vertebrates and *Drosophila*. *Development* **122**: 4023–4031.
- Rodaway, A., H. Takeda, S. Koshida, J. Broadbent, B. Price, J.C. Smith, R. Patient, and N. Holder. 1999. Induction of the mesendoderm in the zebrafish germ ring by yolk cell-derived TGF β family signals and discrimination of mesoderm and endoderm by FGF. *Development* **126**: 3067–3078.
- Saga, Y., S. Miyagawa-Tomita, A. Takagi, S. Kitajima, J. Miyazaki, and T. Inoue. 1999. MesP1 is expressed in the heart precursor cells and required for the formation of a single heart tube. *Development* **126**: 3437–3447.
- Schier, A.F., S.C. Neuhauss, K.A. Helde, W.S. Talbot, and W. Driever. 1997. The *one-eyed p inhead* gene functions in mesoderm and endoderm formation in zebrafish and interacts with *no tail*. *Development* **124**: 327–342.
- Schott, J.J., D.W. Benson, C.T. Basson, W. Pease, G.M. Silberbach, J.P. Moak, B.J. Maron, C.E. Seidman, and J.G. Seidman. 1998. Congenital heart disease caused by mutations in the transcription factor NKX2-5. *Science* **281**: 108–111.
- Schultheiss, T.M., J.B. Burch, and A.B. Lassar. 1997. A role for bone morphogenetic proteins in the induction of cardiac myogenesis. *Genes & Dev.* **11**: 451–462.
- Searcy, R.D., E.B. Vincent, C.M. Liberatore, and K.E. Yutzey. 1998. A GATA-dependent *nkx-2.5* regulatory element activates early cardiac gene expression in transgenic mice. *Development* **125**: 4461–4470.
- Sepulveda, J.L., N. Belaguli, V. Nigam, C.Y. Chen, M. Nemer, and R.J. Schwartz. 1998. GATA-4 and Nkx-2.5 coactivate Nkx-2 DNA binding targets: Role for regulating early cardiac gene expression. *Mol. Cell. Biol.* **18**: 3405–3415.
- Shimoda, N., E. Knapik, J. Ziniti, S. Kaplan, C. Sim, E. Yamada, and M.C. Fishman. 1998. Survey of CA repeats in the zebrafish genome. In *Zebrafish development and genetics*, pp. 274. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Stainier, D.Y.R. and M.C. Fishman. 1992. Patterning the zebrafish heart tube: Acquisition of anteroposterior polarity. *Dev. Biol.* **153**: 91–101.
- Stainier, D.Y.R., R.K. Lee, and M.C. Fishman. 1993. Cardiovascular development in the zebrafish. I. Myocardial fate map and heart tube formation. *Development* **119**: 31–40.
- Stainier, D.Y.R., B. Fouquet, J.N. Chen, K.S. Warren, B.M. Weinstein, S.E. Meiler, M.A. Mohideen, S.C. Neuhauss, L. Solnica-Krezel, A.F. Schier et al. 1996. Mutations affecting the formation and function of the cardiovascular system in the zebrafish embryo. *Development* **123**: 285–292.
- Strahle, U., P. Blader, and P.W. Ingham. 1996. Expression of *axial* and *sonic hedgehog* in wildtype and midline defective zebrafish embryos. *Int. J. Dev. Biol.* **40**: 929–940.
- Suzuki, E., T. Evans, J. Lowry, L. Truong, D.W. Bell, J.R. Testa, and K. Walsh. 1996. The human GATA-6 gene: Structure, chromosomal location, and regulation of expression by tissue-specific and mitogen-responsive signals. *Genomics* **38**: 283–290.
- Vos, P., R. Hogers, M. Bleeker, M. Reijans, T. van de Lee, M. Hornes, A. Frijters, J. Pot, J. Peleman, M. Kuiper et al. 1995. AFLP: A new technique for DNA fingerprinting. *Nucleic Acids Res.* **23**: 4407–4414.
- Warga, R.M. and C. Nüsslein-Volhard. 1999. Origin and development of the zebrafish endoderm. *Development* **126**: 827–838.
- Westerfield, M. 1995. *The zebrafish book*. University of Oregon Press, Eugene, OR.
- Yelon, D., S.A. Horne, and D.Y.R. Stainier. 1999. Restricted expression of cardiac myosin genes reveals regulated aspects of heart tube assembly in zebrafish. *Dev. Biol.* **214**: 23–37.
- Zhu, J., R.J. Hill, P.J. Heid, M. Fukuyama, A. Sugimoto, J.R. Priess, and J.H. Rothman. 1997. *end-1* encodes an apparent GATA factor that specifies the endoderm precursor in *Caenorhabditis elegans* embryos. *Genes & Dev.* **11**: 2883–2896.
- Zhu, J., T. Fukushige, J.D. McGhee, and J.H. Rothman. 1998. Reprogramming of early embryonic blastomeres into endodermal progenitors by a *Caenorhabditis elegans* GATA factor. *Genes & Dev.* **12**: 3809–3814.